



Claims

1. Process for the preparation of chiral compounds of formula




- 10 wherein X represents S or O, and R represents C₁₋₆-alkyl, C₃₋₈-cycloalkyl, aryl or aralkyl, each aryl or aralkyl being optionally further substituted with one or more C₁₋₄-alkyl groups and/or halogen atoms, which process comprises the asymmetric hydrogenation of compounds of formula




- 20 wherein X and R are as defined above, in the presence of a transition metal complex of a chiral bidentate phosphine ligand and, optionally, a base.

2. The process of claim 1 wherein the chiral bidentate phosphine ligand is a compound of formula



wherein R^2 and R^3 are methyl, ethyl or isopropyl; and wherein R^4 and R^5 are hydrogen or R^4 and R^5 together form an isopropylidenedioxy group.

3. The process of claim 1, wherein the chiral bidentate phosphine ligand is a compound of formula



IV,

and enantiomers

wherein R^6 and R^7 are methoxy or ethoxy or wherein R^6 and R^7 together form a 1,3-propylidenedioxy or a 1,4-butyldenedioxy group.

4. The process of claim 1, wherein the chiral bidentate phosphine ligand is selected from the group consisting of (*S,S*)-Me-DuPhos, (*S,S*)-Et-DuPhos, (*S,S,S,S*)-Me-KetalPhos and (*S*)-C4-TunaPhos.
5. The process of any one of claims 1 to 4, wherein the transition metal is Ru or Rh.
6. The process of any one of claims 1 to 5, wherein the transition metal complex of a chiral bidentate phosphine ligand comprises at least one diene, alkene or arene as stabilizing ligand.
7. The process of claim 6, wherein the transition metal complex of a chiral bidentate phosphine ligand comprises at least one stabilizing ligand selected from the group consisting of 1,5-cyclooctadiene and *p*-cymene.
8. The process of any one of claims 1 to 7, wherein the counterion of the transition metal complex of a chiral bidentate phosphine ligand is selected from the group consisting of Cl⁻, BF₄⁻, AsF₆⁻, SbF₆⁻ and triflate.

9. The process of any one of claims 1 to 8, wherein the catalyst is prepared by mixing a transition metal complex of the formula $[\text{Rh}(\text{cod})_2]^+\text{BF}_4^-$ with a chiral bidentate phosphine selected from the group consisting of (*S,S*)-Me-DuPhos, (*S,S*)-Et-DuPhos and (*S,S,S,S*)-Me-KetalPhos.

5


10. The process of any one of claims 1 to 9, wherein the base is a hydroxide, methanolate or ethanolate of lithium, sodium or potassium or a mixture of said bases.

11. The process of any of claims 1 to 10, wherein the hydrogen pressure during the reaction is in the range of 1 to 60 bar and more particularly preferred in the range of 10 to 30 bar.

10

12. Compounds of formula

15



I

and enantiomers

and its addition salts of proton acids, wherein X represents S or O, and R represent C_{1-6} -alkyl, C_{3-8} -cycloalkyl or benzyl with the exception of compounds wherein X is S and R is methyl.

20